arXiv:1406.2213v2 [math.AG] 22 Jun 2014

SOME CASES ON STRASSEN ADDITIVE CONJECTURE

YOUNGHO WOO

Abstract. In this article, we verify the additivity for rank of a sum of coprime monomials and bivariate polynomials generalizing the result in ([CCG12]). We also show similar results hold for cactus rank.

1. Introduction

Given a homogeneous polynimial $F \in S^dV$ of degree d, the *rank* (or *Waring rank*) $\mathrm{rk}(F)$ of F is the minimal number of linear forms $L_1, ..., L_r \in V$ satisfying $F = L_1^d + \cdots + L_r^d$. In [CCG12], the authors show that rank of coprime monomials is the sum of ranks of such monomials. This work verified a particular case of the symmetric version of Strassen additive conjecture. (See [CCC14] and reference therein)

Conjecture 1.1. *If $F \in \mathbb{C}[x_0, ..., x_n]$ and $G \in C[y_0, ..., y_m]$ are homogeneous polynomials of same degree ≥ 2. Then* $\mathrm{rk}(F + G) = \mathrm{rk}(F) + \mathrm{rk}(G)$

In this paper we verify this conjecture for F and G are sums of several coprime monomials and bivariate polymonimals. (See Corollary 3.6 for the precise statement). In the proof, we follows the same track as in [CCG12]. This is possible mainly because for a monomial or a bivariate form, the rank attains the lower bound of the inequality (2) in section 2. In fact, we prove the conjecture for those polynomials satisfying this property. A similar method leads us to the same additivity property for *cactus* rank. (See section 3 and Corollary 4.6)

2. Preliminaries

2.1. **Apolarity.** Let V be a finite dimensional vector space over the complex field $\mathbb{C}$ and $S = SV = \bigoplus_d S^dV$ be the polynomial algebra over $\mathbb{C}$ where S^dV is the symmetric d-product. Similarly, let $T = SV^* = \bigoplus_d S^d(V^*)$ be the polynomial algebra of the dual space V^*. Then a polynomial $g \in T$ acts on S as a linear differential operator. Given $F \in S^dV$, the *perp ideal* $F^\perp \subset T$ is the ideal of elements annihilating F. *i.e.*

$$F^\perp = \{g \in T | g \cdot F = 0\}.$$

It is well-known that the ideal $F^\perp$ is a homogeneous ideal such that the quotient $T/F^\perp$ is a zero-dimensional, Gorenstein algebra with socle degree d. A subscheme $X \subset \mathbb{P}(V)$ is called *apolar* to F if $I_X \subset F^\perp$ where I_X is the (saturated) homogeneous ideal of the subscheme X. Here we think of T as the homogeneous coordinate ring of $\mathbb{P}(V)$.

Now suppose that $F \in S^dV$. Consider, up to scalar, d-th power L^d of a linear form $L \in V$ as an image of the Veronese map $\nu_d : \mathbb{P}(V) \to \mathbb{P}(S^dV)$ defined by

$\nu_d([L]) = [L^d]$. Then the point $[F] \in \mathbb{P}(S^dV)$ is contained in the linear span of the image $\nu_d(\{L_1, ..., L_r\})$ if and only if $F = c_1L_1^d + \cdots + c_rL_r^d$ for some $c_1, ..., c_r$. Therefore the rank of F is

$$\min\{r|[F] \in < \nu_d(X) >, X \subset \mathbb{P}(V) \text{ is reduced, } \dim(X) = 0 \text{ and} \deg(X) = r\}$$

, where $< \cdot >$ means its linear span in $\mathbb{P}(S^dV)$.
The *cactus* rank of F (denoted by $\mathrm{crk}(F)$) is defined similarly,

$$\mathrm{crk}(F) = \min\{r|[F] \in < \nu_d(X) >, \ \dim(X) = 0 \text{ and} \deg(X) = r\}.$$

Note that in this case, we drop the condition "reduced"and consider all 0-dimensional subschemas $X \subset \mathbb{P}(V)$ of degree r.

Lemma 2.1 (Apolarity Lemma [IK99])**.** *Let $F \in S^dV$ and $X \subset \mathbb{P}(V)$ be a zero dimensional subscheme. The point $[F] \in \mathbb{P}(S^dV)$ is in the linear span of $\nu_d(X)$ if and only if the subscheme X is apolar to F. i.e. $I_X \subset F^\perp \subset T$.*

Remark 2.2. Let $F \in S^dV$. According to the apolarity lemma, we can reformulate the definition of rank and cactus rank

(1) $\mathrm{rk}(F) = \min\{\deg(X)|X \subset \mathbb{P}(V)$ is 0-dimensional, reduced and $I_X \subset F^\perp\}$
(2) $\mathrm{crk}(F) = \min\{\deg(X)|X \subset \mathbb{P}(V)$ is 0-dimensional and $I_X \subset F^\perp\}$

2.2. **Degree of a zero-dimensional subscheme.** Let $X \subset \mathbb{P}(V)$ be a zero-dimensional scheme, and let $I_X \subset T$ be the homogeneous ideal of X. Then for a linear form $l \in V^*$ such that $X \cap \{l = 0\} = \phi$, we have

$$\deg(X) = \dim_{\mathbb{C}} T/(I_X + (l)) = \sum_{i=0}^{\infty} H(T/(I_X + (l)), i),$$

where $H(T/(I_X + (l)), \cdot)$ is the Hilbert function of $T/(I_X + (l)$.
For a given $F \in S^dV$, if X is apolar to F, then we have $\deg(X) \geq \dim_{\mathbb{C}} T/(F^\perp + (l))$. On the other hand, the ideal $F^\perp + (l)$ defines a zero-dimensional subscheme in $\mathrm{Spec}(T)$. The length of this subscheme attains the minimum for a general l, by upper semi-continuity. So we conclude that

$$\mathrm{crk}(F) \geq \dim_{\mathbb{C}} T/(F^\perp + (l)) \text{ for a } \textit{general} \text{ linear form } l \in V^*. \tag{1}$$

Now $t \in V^*$ be a linear form, assume that X is reduced. Then the ideal quotient $(I_X : t)$ is the homogeneous ideal of the reduced subscheme $X \setminus \{t = 0\}$. As above, if X is apolar to F, then we have $\deg(X) \geq \dim_{\mathbb{C}} T/((F^\perp : t) + (t))$ and

$$\mathrm{rk}(F) \geq \dim_{\mathbb{C}} T/((F^\perp : t) + (t)) \text{ for } \textit{any} \text{ linear form } t \in V^*. \tag{2}$$

3. Rank

Definition 3.1. Let $F \in S^dV$ be a homogeneous polynomial of degree d, and $t \in V^*$. We will say that the rank of F is *computed by the linear form* t if

$$\mathrm{rk}(F) = \dim_{\mathbb{C}} T/(F^\perp : t) + (t).$$

Example 3.2 (**Bivariate case**)**.** Suppose that $\dim V = 2$. Then for every $F \in S^dV$, the rank of F is computed by some linear form $t \in V^*$. In fact, it is well-known that $F^\perp = (g_1, g_2)$ for some homogeneous polynomials $g_1, g_2 \in T$ of degree $d_1 \leq d_2$ respectively. Let t be any linear factor of g_1. Then the ideal quotient

$F^\perp : t = (g_1/t, g_2)$. In fact, it is obvious that $(g_1/t, g_2) \subset F^\perp : t$. Using an exact sequence

$$0 \to T/(F^\perp : t) \to T/F^\perp \to T/(F^\perp, t) \to 0,$$

we have $\dim T/(F^\perp : t) = (d_1 - 1)d_2 = \dim T/(g_1/t, g_2)$ because t is not a factor of g_2. Hence $F^\perp : t = (g_1/t, g_2)$ holds. Now we have two separate cases

(1) (g_1 has a double root) Then $\mathrm{rk}(F) = d_2$. Let t be the multiple factor of g_1. Since g_1/t still contains t, $\dim_{\mathbb{C}} T/(F^\perp : t) + (t) = d_2$
(2) (g_1 has no double roots) Then $\mathrm{rk}(F) = d_1$. If $d_1 < d_2$, then take t as one of linear factors of g_2 and $\dim_{\mathbb{C}} T/(F^\perp : t) + (t) = d_1$. If $d_1 = d_2$ and g_2 has no double roots, then we can choose $g_1' = c_1 g_1 + c_2 g_2$ for some constant c_1, c_2 such that g_1' has a double root and $F^\perp = (g_1', g_2)$ which belongs to the firtst case.

Example 3.3 (**Monomial cases**). ([CCG12], [BBZ13]) Let $F = x_0^{a_0} \cdots x_1^{a_n} \in \mathbb{C}[x_0, ..., x_n]$. Let $\{t_0, ..., t_n\}$ be the dual basis of $\{x_0, ..., x_n\}$. Assume that $a_0 \leq a_i$ for all i. Then $F^\perp = (t_0^{a_0+1}, ..., t_n^{a_n+1})$ and $\mathrm{rk}(F) = (a_1 + 1) \cdots (a_n + 1)$. Easy to see that $(F^\perp : t_0) + (t_0) = (t_0, t_1^{a_1+1}, ..., t_n^{a_n+1})$. Therefore $\mathrm{rk}(F) = \dim T/(F^\perp : t_0) + (t_0))$ is computed by t_0.

Now we fix some notations. Suppose that we have a decomposition $V = V_1 \oplus V_2 \oplus \cdots \oplus V_m$ for some m and the dual decomposition $V^* = V_1^* \oplus V_2^* \oplus \cdots \oplus V_m^*$. Then there is an natural idenfication $S(V_i)$ to a subspace of $S = SV$ (and $S(V_i^*)$ to SV^* resp.) for each i. By abuse of notation, we will set $S_i = S(V_i)$ and $T_i = S(V_i^*)$. Suppose that $F_i \in S^d(V_i)$ and $t_i \in V_i^*$ for $i = 1, ..., m$, and let $F = F_1 + \cdots + F_m \in S^d V$. Fixing such F_i and t_i for all i, we put $J_i = (F_i^\perp : t_i) + (t_i, V_1^*, ..., V_{i-1}^*, V_{i+1}^*, ..., V_m^*)$. **Here $F_i^\perp$ means the perp ideal of F_i *in* T_i.** For instance, J_1 is the ideal generated by

- $g \in T_1$ with $t_1 g \in F_1^\perp$,
- t_1 and
- all linear forms in $V_2^*, , ..., V_m^*$.

under the inclusion $T_i \subset T$.

Lemma 3.4. *Notation as above.* $(F^\perp : (t_1, ..., t_m)) \subset J_1 \cap \cdots \cap J_m$.

PROOF. By symmetry, it is enough to show that $F^\perp : t_1 \subset (F_1^\perp : t_1) + (t_1, V_2^*, ..., V_m^*)$. Let $h \in (F^\perp : t_1)$. We can write $h = h_1 + h_2$ where $h_1 \in T_1$ and $h_2 \in (V_2^*, ..., V_m^*)$. We claim that $h_1 \in (F_1^\perp : t_1)$. By definition, $t_1 h \in F^\perp$. Since $t_1 h_2 \in F^\perp$, we have $t_1 h_1 = t_1 h - t_1 h_2 \in F^\perp \cap T_1 = F_1^\perp$ and hence $h_1 \in (F_1^\perp : t_1)$. The inclusion follows.

Proposition 3.5. *Suppose that we have a decompsition $V = V_1 \oplus V_2 \oplus \cdots \oplus V_m$ for some m. Let $F_i \in S^d V_i$ for $1 \leq i \leq m$ and for $d \geq 2$, let $F = F_1 + ... + F_m \in S^d V$. If the rank of F_i is computed by some $t_i \in V_i^*$ for all i, then* $\mathrm{rk}(F) = \mathrm{rk}(F_1) + ... + \mathrm{rk}(F_m)$.

The proof proceeds similarly as in [CCG12]
PROOF. We will prove that

$$\mathrm{rk}(F) \geq \mathrm{rk}(F_1) + \cdots + rk(F_m) \tag{3}$$

because the opposite inequality is obvious.
Let $r = \mathrm{rk}(F)$. Then by apolarity lemma, there is a finite set $X \subset \mathbb{P}(V)$ of degree r such that the (saturated) homogeneous ideal $I_X \subset F^{\perp}$. Then the quotient ideal $I_X : (t_1, t_2, .., t_m)$ is the homogeneous ideal of the subset X' of X not lying on $\{t_1 = t_2 = \cdots = t_m = 0\}$. Then a linear form $\lambda_1 t_1 + \cdots \lambda_m t_m \in V^*$ for general $\lambda_1, ..., \lambda_m \in \mathbb{C}$ does not vanish any point of X'. By replacing t_i by $\lambda_i t_i$, we may assume that $X' \cap \{t_1 + \cdots + t_m = 0\} = \phi$. By the previous lemma and the fact $(t_1 + \cdots + t_m) \subset J_1 \cap \cdots \cap J_m$, we have

$$I_{X'} + (t_1 + \cdots + t_m) \subset J_1 \cap \cdots \cap J_m \tag{4}$$

Now we prove two claims.

Claim 1. $\dim_{\mathbb{C}} T/(J_1 \cap \cdots \cap J_m) = \mathrm{rk}(F_1) + \cdots \mathrm{rk}(F_m) - (m-1)$.
For each $i = 1, ..., 2$, one has $T/J_i \cong T_i/((F_i^{\perp} : t_i) + (t_i))$, and hence $\dim T/J_i = \mathrm{rk}(F_i)$. From the exact sequence

$$0 \to T/(J_1 \cap \cdots \cap J_m) \to T/(J_1 \cap \cdots \cap J_{m-1}) \oplus T/J_m \to T/(J_1 \cap \cdots \cap J_{m-1} + J_m) \to 0,$$

and $T/(J_1 \cap \cdots \cap J_{m-1} + J_m) \cong \mathbb{C}$, the assertion follows by induction.

Claim 2. $t_1, ..., t_m$ are linearly independent modulo $I_{X'}$.
Suppose $\mu_i t_1 + \cdots + \mu_m t_m \in I_{X'}$ for some $\mu_1, ..., \mu_m \in \mathbb{C}$. Since this already vanishes along $\{t_1 = \cdots = t_m = 0\}$, we have $\mu_1 t_1 + \cdots \mu_m t_m \in I_X \subset F^{\perp}$. Then $(\mu_1 t_1 + \cdots \mu_m t_m) \cdot F = \mu_1 t_1 \cdot F_1 + \cdots + \mu_m t_m \cdot F_m = 0$. Note that each polynomial $t_i \cdot F_i \in S_i (= S(V_i))$ has degree ≥ 1 or $t_i \cdot F_i = 0$ for $i = 1, ..., m$. Since t_i compute the rank of F_i, $t_i \cdot F_i \neq 0$. (Otherwise $1 \in (F_i : t_i)$.) Thus $\mu_1 = \cdots = \mu_m = 0$

From the claim 2 and the inclusion (4), we conclude that

$$H(T/(I_{X'} + (t_1 + \cdots + t_m)), 1) \geq H(T/(J_1 \cap \cdots \cap J_m, 1) + (m-1) \tag{5}$$

since $t_1, ..., t_m \in J_1 \cap \cdots \cap J_m$.
Now we verify the inequality (3).

$$\begin{aligned}
r \geq & \deg(X') = \dim T/((I_{X'} + (t_1 + \cdots + t_m)) \\
= & \sum_{j=0}^{\infty} H(T/(I_{X'} + (t_1 + \cdots + t_m)), j) \\
= & H(T/(I_{X'} + (t_1 + \cdots + t_m)), 1) + \sum_{j \neq 1} H(T/(I_{X'} + (t_1 + \cdots + t_m)), j) \\
\geq & H(T/(J_1 \cap \cdots \cap J_m, 1) + (m-1) + \sum_{j \neq 1} H(T/(J_1 \cap \cdots \cap J_m, j) \text{ by (4) and (5)} \\
= & \dim_{\mathbb{C}} T/(J_1 \cap \cdots \cap J_m) + (m-1) \\
= & \mathrm{rk}(F_1) + \cdots \mathrm{rk}(F_m) \text{ by the claim 1}
\end{aligned}$$

Corollary 3.6. *Let $F_i \in S^d V_i$ $(d \geq 2)$ be a monomial or a bi-variate polynomial for each i. Then* $\mathrm{rk}(F_1 + \cdots + F_m) = \mathrm{rk}(F_1) + \cdots + \mathrm{rk}(F_m)$.

4. Cactus Rank

Definition 4.1. Let $F \in S^dV$ be a homogeneous polynomial of degree d. We will say that the cactus rank of F is *computed by a general linear form* if

$$\mathrm{crk}(F) = \dim_{\mathbb{C}} T/(F^{\perp} + (l)).$$

for a general linear form $l \in V^*$.

Example 4.2. Suppose that $\dim V = 2$. Then for every $F \in S^dV$, the cactus rank of F is computed by a general linear form. As in the example (3.2), $F^{\perp} = (g_1, g_2)$ for some homogeneous polynomials $g_1, g_2 \in T$ of degree $d_1 \leq d_2$ respectively. Then $\mathrm{crk}(F) = d_1 = \dim_{\mathbb{C}} T/((g_1, g_2) + (l))$ for a general $l \in V^*$.

Example 4.3. ([RS11]) As in the example (3.3), let $F = x_0^{a_0} \cdots x_1^{a_n} \in \mathbb{C}[x_0, ..., x_n]$. Here we give an additional assumption that $a_0 + \cdots + a_{n-1} \leq a_n$. Then $F^{\perp} = (t_0^{a_0+1}, ..., t_n^{a_n+1})$ and $\mathrm{crk}(F) = (a_0 + 1) \cdots (a_{n-1} + 1)$ is computed by a general linear form. In fact, for general $\lambda_0, ..., \lambda_n \in \mathbb{C}$

$$\begin{aligned}
&\dim_{\mathbb{C}} T/(F^{\perp} + (\lambda_0 t_0 + \cdots \lambda_n t_n)) \\
&\quad = \dim_{\mathbb{C}} \mathbb{C}[x_0, ..., x_{n-1}]/(t_0^{a_0+1}, ..., t_{n-1}^{a_{n-1}+1}, (t_0 + \cdots + t_{n-1})^{a_n+1}) \\
&\quad = \dim_{\mathbb{C}} \mathbb{C}[x_0, ..., x_{n-1}]/(t_0^{a_0+1}, ..., t_{n-1}^{a_{n-1}+1}) \\
&\quad = (a_0 + 1) \cdots (a_{n-1} + 1)
\end{aligned}$$

because $(t_0 + \cdots + t_{n-1})^{a_n+1} \in (t_0^{a_0+1}, ..., t_{n-1}^{a_{n-1}+1})$ provided that $a_0 + \cdots + a_{n-1} \leq a_n$

From now on, fix $F \in S^dV$ for $d \geq 2$. Suppose that $F = F_1 + F_2$ for some $F_1 \in S^dV_1$ and $F_2 \in S^dV_2$ under the decomposition $V = V_1 \oplus V_2$.

Lemma 4.4. *Let $l_i \in V_i^*$ for $i = 1, 2$. Suppose that $d \geq 2$ and $l_i \notin F_i^{\perp}$. Then*

(1) $F^{\perp} + (l_1 + l_2) \subsetneq [(F_1^{\perp} + (l_1) + (V_2^*)] \cap [F_2^{\perp} + (l_2) + (V_1^*)]$
(2) $\dim T/(F^{\perp} + (l_1 + l_2)) \geq \dim T_1/(F_1^{\perp} + (l_1)) + \dim T_2/(F_2^{\perp} + (l_2))$

PROOF. At first, we show that $F^{\perp} \subset F_1^{\perp} + (l_1) + (V_2^*)$. Let $h \in F^{\perp}$ be a nonzero homogenous polynomial. Since the set $F_1^{\perp} + (V_2^*)$ contains all monomials of degree $> d$, we may assumt that $\deg(h) \leq d$. Now we can write $h = h_1 + h_2$ for $h_1 \in T_1$ and $h_2 \in V_2^*$. Since $h \cdot F = 0$, we have $h_1 \cdot F_1 + h_2 \cdot F_2 = 0$. If $\deg(h) < d$, then $h_1 \cdot F_1 = h_2 \cdot F_2 = 0$ and hence $h \in F_1^{\perp} + (V_2^*)$. On the other hand, if $\deg(h) = d$, then both $h_1 \cdot F_1$ and $h_2 \cdot F_2$ are constant. Since $l_1 \notin F_1^{\perp}$, there is $g \in T_1$ of degree $d - 1$ with $g \cdot (l_1 \cdot F_1)$ is a nonzero constant. Then $h_1 - \frac{(h_1 \cdot F)}{g l_1 \cdot F_1} g l_1 \in F_1^{\perp}$. So we have $h = (h_1 - \frac{(h_1 \cdot F)}{g l_1 \cdot F_1} g l_1) + \frac{(h_1 \cdot F)}{g l_1 \cdot F_1} g l_1 + h_2 \in F_1^{\perp} + (l_1) + (V_2^*)$. Up to symmetry, the inclusion of the lemma follows. This inclusion must be strict because $l_1 \notin F^{\perp} + (l_1 + l_2)$. Otherwise, there is a constant λ such that $(1 - \lambda) l_1 - \lambda l_2 \in F^{\perp}$, and hence $(1 - \lambda) l_1 \cdot F_1 - \lambda l_2 \cdot F_2 = 0$. But this is impossible since $\deg(F) \geq 2$ and $l_i \notin F_i^{\perp}$ for $i = 1, 2$. This proves (1).
Now Put $J_1 = F_1^{\perp} + (l_1) + (V_2^*)$, and $J_2 = F_2^{\perp} + (l_2) + (V_1^*)$ as in the rank case.

By (1), we have

$$\begin{aligned}
&\dim T/(F^\perp + (l_1 + l_2)) \\
>&\dim T/J_1 \cap J_2 \\
=&\dim T/J_1 + \dim T/J_2 - \dim T/(J_1 + J_2) \\
=&\dim T_1/(F_1^\perp + (l_1)) + T_2/(F_2^\perp + (l_2)) - 1
\end{aligned}$$

Proposition 4.5. *Suppose that the cactus rank of F_i is computed by a general linear form for $i = 1, 2$. Then* $\mathrm{crk}(F) = \mathrm{crk}(F_1) + \mathrm{crk}(F_2)$ *and this is computed by a general linear form.*

PROOF. We always have $\mathrm{crk}(F) \leq \mathrm{crk}(F_1) + \mathrm{crk}(F_2)$. By assumption, $\mathrm{crk}(F_i) = \dim T_i/(F_i^\perp + (l_i))$ for a general linear form l_i for $i = 1, 2$. Then $l_i \notin F_i^\perp$. By apolarity lemma, there is a zero dimensional subscheme $X \subset \mathbb{P}(V)$ such that $I_X \subset F^\perp$ and $\deg(X) = \mathrm{crk}(F)$. Since $l_1 + l_2$ is a general linear form in V^*, we have

$$\begin{aligned}
\deg(X) =& \dim T/(I_X + (l_1 + l_2)) \\
\geq& \dim T/(F^\perp + (l_1 + l_2)) \\
\geq& \dim T_1/(F_1^\perp + (l_1)) + \dim T_2/(F_2^\perp + (l_2)) \qquad \text{(by the lemma)} \\
=& \mathrm{crk}(F_1) + \mathrm{crk}(F_2)
\end{aligned}$$

Corollary 4.6. *Let $F_i \in S^d V_i$ $(d \geq 2)$ be a monomial satisfying the assumption in example 4.3 or a bi-variate polynomial for each $i = 1, ..., m$. Then* $\mathrm{crk}(F_1 + \cdots + F_m) = \mathrm{crk}(F_1) + \cdots + \mathrm{crk}(F_m)$.

References

[AH95] J. Alexander and A. Hirschowitz. Polynomial interpolation in several variables. *J. Algebraic Geom.*, 4(2):201–222, 1995.

[BBZ13] W. Buczynska, J. Buczynski, Z. Teitler. Waring decompositions of monomials. *J. Algebra* 378 (2013), 4557.

[BR13] A. Bernardi, K. Ranestad. On the cactus rank of cubics forms. *J. Symbolic Comput.* 50: 291–297, 2013

[CCC14] Enrico Carlini, Maria Virginia Catalisano, Luca Chiantini. Progress on the symmetric Strassen conjecture. *arXiv:1405.3721*

[CCG12] E. Carlini, M.V. Catalisano, and A.V. Geramita. The solution to the Waring problem for monomials and the sum of coprime monomials. *J. Algebra*, 370:5–14, 2012.

[IK99] A. Iarrobino and V. Kanev. Power sums, Gorenstein algebras, and determinantal loci, volume 1721 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1999.

[Lan12] J.M. Landsberg. Tensors: geometry and applications, volume 128 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2012.

[LT10] J.M. Landsberg, Z. Teitler, On the ranks and border ranks of symmetric tensors, *Found. Comput. Math.* 10 (3) (2010) 339366. MR2628829 (2011d:14095)

[RS11] K. Ranestad, F. Schreyer, On the rank of a symmetric form. *J. Algebra* 346 : 340–342, 2011

[Str73] V. Strassen. Vermeidung von Divisionen. *J. Reine Angew. Math.*, 264:184–202, 1973.

Department of Mathematical Sciences, KAIST, Taejeon, South Korea
E-mail address: youngw@kaist.ac.kr